

# Projecting Births in Iran Using a Three-Parameter Model

Meimanat Hosseini and Peter McDonald

Demography Unit

Melbourne School of Population and Global Health

University of Melbourne

# Background

- In a previous paper based on Australia, the argument was made that, for all births beyond the first birth, parity and interval since the last birth are much better predictors of the number and timing of future births than is age – particularly age on its own.
- McDonald, P. and Kippen , R. 2011. Forecasting births. Feature Article, ABS Catalogue Number 2051.0, Australian Bureau of Statistics, Canberra.  
<http://www.abs.gov.au/ausstats/abs@.nsf/mf/2051.0>

# Rationale: Probabilities

- The patterning of births is such that parity is the principal determinant of whether or not a woman will have another birth and that time since the last birth is a good predictor of when she will have the next birth.
  - A 37-years-old women who had her first birth three years ago is much more likely to have the next birth than a 27 year-old woman who had her first birth 10 years ago.

# **Rationale: Population at risk**

- McDonald and Kippen demonstrated in the Australian case that there was a high degree of stability across time in age-parity-time since last birth probabilities of birth for all births other than the first. The principal determinant of change then became the rate and timing of first births.
- Changes in the timing of first births produced changes in the relevant population at risk by age, parity and time since last birth and forecasting accuracy was greatly enhanced by taking the three-parameter population into consideration because it took into account previous changes in first births.

# The centrality of the first birth

The Australian study concluded:

When fertility rates are low, the main factor that affects the **annual** incidence of births is changes in the rates of first birth by age across successive cohorts of women.

# Data requirements: two large matrices

- 1. A matrix of probabilities of birth by single years of age (35), single parity (10) and single years since the time of the last birth (10) = 3500 cells (but a large number of the cells are zero or near zero).
- 2. An equivalent (3500 cell) matrix of the population at risk (but again a large number of the cells are zero or near zero).
- Deriving the database requires a very large survey or census, or accumulation from birth registration data.

# **Data inputs for Iran**

---

## **1) 1986, 1996, 2006 Censuses**


## **2) The 2000 Iran Demographic Health Survey (IDHS)**

Households interviewed:	<b>113,913</b>
Ever married women 15-49:	<b>90,240</b>
Births occurred to women:	<b>316,371</b>

## **3) The 2010 Iran Multiple Indexes & Demographic Health Survey (IDHS)**

Households interviewed:	<b>31,350</b>
Ever married women 15-54:	<b>34,435</b>
Births occurred to women:	<b>62,189</b>


**Figure 18. Total Fertility Rate, Iran, 1980-2009**




Data sources: The 2010 and 2000 Iran DHS and the 2006 and 2011 Censuses





## Proportion Never Married by Age, Real Birth Cohorts





# Probability of first birth by years since marriage for women's age at marriage





# Probability of having first birth by years since marriage for selected years in which the first birth occurred



# Probability of having second birth by years since first birth for selected years in which the second birth occurred




# Probability of having third birth by years since second birth for selected years in which the third birth occurred






## Second births


2000




2010



## Third births




Years since previous parity: 0 1 2 3 4 5 6 7 8 9 10+



Years since previous parity: 0 1 2 3 4 5 6 7 8 9 10+



# Number of registered births and forecasted births in the equivalent years between 2000 and 2010



# Limitations

- Getting the data, especially the population at risk.
- Smoothing the data surface for random fluctuations where small numbers are involved.
- Handling the data. Difficult with Excel because of the size. Need to use matrix algebra and a computer program.
- Method for projecting probabilities of first birth.